

Title: "PROCESS AND COMPOSITION FOR ENHANCING THE ACTION OF VITAMIN A
ON THE CELLULAR ACTIVITY OF AN INDIVIDUAL, AND USE OF VITAMIN C"

5 **Field of the Invention**

The present invention refers to a process for improving the effects of Vitamin A used in cosmetic compositions in order to enhance the cellular activity of an individual.

Background of the Invention

10 The compound generically known as Vitamin A comprises retinol and its derivatives, also known as retinoids, in addition to its acidic or aldehyde form, respectively retinoic acid and retinal. Retinoic acid has application in the pharmaceutical and cosmetic industries being, however, prohibited in several countries for cosmetic use due to the adverse effects of irritability which it may cause. Examples of pharmaceutical applications of retinoic acid can be found in the article "Relationships between structure and activity of retinoids", published
15 by Nature, Volume 236, pages 110-113, of September 9, 1996.

In the cosmetic area Vitamin A is usually employed in the form of retinol or some of its retinoids such as retinyl palmitate, and the use of retinol causes various biologic activities, many of which are highly desirable in cosmetic compositions, particularly in those intended to improve the general conditions of the skin of the individual subjected to the topic
20 use thereof. Results achieved by the topic use of Vitamin A are described in passages contained in pages 82 - 119 of the article entitled "Vitamin A Complex", written by Wade Cheng, PhD and Shirley DePetris and published by Skin Inc., March/April 1998.

Moreover, regulation and balance of the epidermal cellular growth through the total synthesis of collagen, among others, such as retention of water in the skin, are also
25 known as effects resulting from the use of Vitamin A in its pure form, called Retinol.

One problem resulting from the use of Vitamin A, either in its pure form or as a derivative, is that, on the one side it promotes the effects of increasing the cellular activity at the level of the dermis and epidermis, accelerating the process of proliferation and differentiation of the keratinocytes and reorganization of the fibers of the dermis (collagen and elastin). But on the other side it must be administered at low doses due to its toxicity. This fact limits the use of Vitamin A and its derivatives to lower contents or requires the utilization of other means that are able to minimize the discomfort of irritation in the skin.

In fact, the use of retinol at low contents is quite common, as shown by several studies, such as the one conducted by the Hamburg Clinic of Experimental Dermatology, in Germany, which discloses tests with low contents of Retinol (0.034%) for men and women with age between 22 and 34 years and which show that such a concentration of retinol could reduce the amount and the deepness of wrinkles. Therefore, this study generically shows the effect of reducing wrinkles by the use of low contents of retinol.

On the other hand, what has been observed is that, even though low concentrations of retinol effectively cause little or no irritation, the results on the skin can remain below the desired levels for the present standards of demand of the consumers in view of the small amount of retinol incorporated in the cosmetic composition and available for its biological action.

In this respect, there have been attempts to obtain compositions of Vitamin A that present effective action and do not cause adverse effects, for instance, the irritation of the skin. As an example, documents US 5516793 and US 5703122 in the name of Avon Products, Inc., are incorporated herein as prior art references. These documents describe a generic association of amounts ranging from 0.5 to 25% by weight of Vitamin C with several irritating active principles, among which Vitamin A is included. This association, however, has the exclusive purpose of reducing irritation of the skin caused by Vitamin A.

It is therefore an objective of the present invention to provide an alternative for the use of Vitamin A at such concentrations that enable an increase in its properties which are beneficial to the skin, without presenting the problems cited above.

Summary of the Invention

The present invention refers to a process for enhancing the action of Vitamin A on the cellular activity of an individual comprising the association of Vitamin C with Vitamin

A, which will be applied to the referred-to individual at a weight ratio ranging from about 1:1 to about 10:1.

In another aspect, the invention refers to a composition for enhancing the action of Vitamin A on the cellular activity of an individual comprising Vitamin C in association with Vitamin A at a weight ratio in the range from about 1:1 to 10:1.

The invention further refers to the use of Vitamin C for enhancing the action of Vitamin A on the cellular activity of an individual.

Brief Description of the Drawings

- Figure 1 shows a graph representing the increase obtained in the cellular activity of a reconstituted skin by the synergistic effect of an association of Vitamin C with Vitamin A according to the invention as compared to the cellular activity of a reconstituted skin treated only with pure Vitamin A.

- Figure 2 shows the synergistic effect on the recuperation and the increase in the cellular activity in reconstituted skin treated with Vitamin A associated with Vitamin C when subjected to ultraviolet irradiation.

Detailed Description of the Invention

After detailed studies the inventors have found that the association of Vitamin C added to compositions containing Vitamin A at a weight ratio ranging from about 1:1 to about 10:1, preferably from about 1:1 to about 5:1, and more preferably from about 1:1 to about 2:1, provides a surprising increase in the cellular activity effects of Vitamin A on an individual.

"Vitamin C" useful for the present invention comprises Vitamin C in its pure form or its derivatives, namely L-ascorbic acid in its molecular form as well as its salts and esters such as ascorbyl phosphate.

As used herein, the expression "an increase in the cellular activity" means the occurrence of a benefit brought about by the increase or improvement at least in one of the situations selected from the maintenance of the cellular condition, the cellular proliferation and the metabolic activity especially in cutaneous cells.

Tests carried out on reconstituted skin show that a treatment of the skin *in vi-*
tro with the association of Vitamin C with Vitamin A according to the present invention pro-
motes an unexpected synergistic increase in the cellular activity of 100% as compared to the
cellular activity observed in the same skin treated exclusively with pure Vitamin A or retinol
5 (figure 1).

In the same surprising way, it has been noted that the association of Vitamin
C with Vitamin A promotes reconstitution, recuperation and increase in the cellular activity of
the skin, even when the individual is subjected to ultraviolet irradiation, which is recognized
to cause deleterious effects on the skin and its cells. Tests carried out to this respect show a
10 synergistic effect of reconstitution and increase in the cellular activity of 5% on reconstituted
skin treated with Vitamin A associated with Vitamin C when subjected to ultraviolet irradiation
(figure 2).

The association of Vitamin C with Vitamin A according to the present invention
may be carried out at the moment of the application of these compounds to the individual,
15 but it can also be advantageously formulated as a cosmetic composition containing the two
vitamins at a weight ratio ranging from about 1:1 to about 10:1, preferably from about 1:1 to
about 5:1, and more preferably from about 1:1 to about 2:1 of Vitamin C to Vitamin A.

According to a preferred embodiment of the invention, said cosmetic compo-
sition comprises, by weight, about 0.01 to about 0.9% of Vitamin C and from about 0.008 to
20 about 0.20% of Vitamin A, based on the total weight of the composition. Even more prefera-
bly, the composition contains from about 0.02 to about 0.8% by weight of Vitamin C and from
about 0.009 to about 0.16% by weight, of Vitamin A and even more preferably the composi-
tion contains 0.02 wt.% of Vitamin C and from about 0.009 to about 0.02 wt% Vitamin A, all
the percentages based on the total weight of the composition.

25 It is noted that, even at very low concentrations, Vitamin A associated with
Vitamin C as defined in the present invention achieves the desired effects of increase in the
cellular activity.

The cosmetic compositions containing Vitamin A and Vitamin C at the propor-
tions cited above can also contain other appropriate additives and formulation aids, such as
30 antioxidants for combating free radicals. Among the useful antioxidants, Vitamin E stands
out, both in its pure form presented by di- α -tocopherol, and as its derivatives such as di- α -
tocopherol, or 2,6-di-terc-buthyl-p-cresol (BHT).

- 5 -

The introduction of Vitamins C, A and E in microspheres increases their action and makes it possible for them to reach the deeper layers of the skin with greater, or even total, integrity, without degeneration of the product in the path between the application area and the place of action.

- 5 In a particularly preferred way, the cosmetic compositions according to the present invention are formulated in such a manner, that their components are contained in organic vectors such as microspheres and, more particularly, in microspheres or microcapsules containing biologically active material ("Talasferas") such as those defined in US Pat 5,395,620, or in Brazilian patent application PI 9706994-7, filed in the name of this same applicant.
- 10

- The composition as described above may contain a plurality of said microspheres, in a dispersed form, comprising Vitamin A and, for example, an antioxidant such as Vitamin E, inserted into a first group of microspheres, and Vitamin C inserted into a second group of microspheres. A particularly preferred composition comprises a first group of microspheres containing Vitamin A at an average concentration of 0.014% and Vitamin E at an average concentration of 0.0005% by weight, and a second group of microspheres containing 0.02% by weight of Vitamin C.
- 15

- Advantageously, in association to the groups of microspheres previously mentioned, such a composition may further contain, in addition to Vitamin A and Vitamin E, cosmetic compounds selected from the group comprising skin structurers, preferably squalan and sphingolipide complexes, skin micronutrients, preferably seaweed extract, sensorial agents, for example, moisteners such as glycerin and hydroxy prolisilan C, emollients such as buthylene glycol and cethyl lactate and silicones such as cyclomethicone, solar protection factors such as Parsol 1789 and Eusolex 6300, emulsifiers, preferably Carbopol 1342 associated to trietanolamin and soybean lecithin, thickeners, preferably xanthan gum; sequestrants, preferably EDTA, antioxidants such as BHT and dl- α -tocopherol, fragrances, conservants, water and mixtures thereof.
- 20
- 25

- In one particular embodiment of the present invention, the composition containing Vitamin A and Vitamin C may be in the form of an emulsion and, in this case, the Vitamin C preferably used is L-ascorbic acid stabilized by hydrogen-bridge-forming compounds. Such processes of stabilizing L-ascorbic acid are described in applications PI 9704418-0 and PI 9704728-7, also filed by this same applicant.
- 30

As an illustrative example of another possible embodiment of the present invention, the composition is formulated as a gel in which the weight ratio of Vitamin C to Vitamin A is advantageously about 5:1, Vitamin C being present preferably in amounts of about 0.75% and Vitamin C being present in amounts of about 0.16 wt.%, based on the total weight of the composition. This gel composition may further contain thickeners such as car-
bopol, fragrances, conservants and water.

THE **NEW** **YORK** **PUBLIC** **LIBRARY**